

Insights on the Depression and Anxiety

Volume - 8, Issue - 1

Review Article **Published Date:-2024-06-13 10:56:40**

[The Primary Care Treatment System for Severe Depression: Perspectives of Patients, Doctors, Treatment Guidelines and Treatment System Failures](#)

There are a number of key issues that matter to patients and General Physicians (GPs) in the primary care treatment system for severe depression. Patient and doctor narratives can contribute by highlighting these key issues. Various systems are employed in treating severe depression. However, there needs to be an investigation using systems failure methodology and how this methodology is applied which can help identify how and why the NHS treatment system for severe depression can fail patients in terms of provision of effective care.

Research Article **Published Date:-2024-04-15 15:19:54**

[Alcohol and Substance Abuse in the General Population during the COVID-19 Pandemic: Results of the COMET-G International Study](#)

The global impact of the COVID-19 pandemic on mental health and substance use behaviors has sparked extensive research efforts. The COMET-G international study, organized by the Department of Medicine and the Rectorate of the Aristotle University of Thessaloniki in collaboration with the World Psychiatric Association, delved into these issues. Running from March 2020 to April 2021, the study collected responses from 55,589 individuals across 40 countries. Through a comprehensive questionnaire, participants provided insights into their mental state, attitudes toward the pandemic, and the resultant changes in their personal and daily lives. Findings revealed, among other things, significant patterns of change in substance use, with notable correlations between reduced usage and the severity of lockdown measures among non-binary individuals. Mental health history emerged as a strong predictor of substance use changes, with influences from anxiety disorders, depression, and self-harm. Additionally, family and social dynamics, including economic expectations and household composition, significantly shaped substance use behaviors during lockdowns. Given these findings, the development of comprehensive approaches targeting the adverse effects of the pandemic on individual behaviors and general welfare is crucial.

Research Article **Published Date:-2024-03-12 16:17:48**

[Death Wishes, Aging Patients, and Euthanasia](#)

The authors are searchers in psychopathology and communicate here about the dead wishes in relation to euthanasia. In Europe, the question comes regularly up to know if the law should be changed concerning the prohibition of euthanasia. The health system obeys progressively a modern idea of comfort and the "good life". The authors are psychotherapists and their methodology is based mainly on phenomenological psychology, psychoanalysis, and psychopathology. Statistics of the French Ministry of Health will support their statements. Different clinical experiences with young patients, aging patients, or near-to-death patients are crossed and compared to those marked by heavy experiences like rape or amputation. The unbearable nature of their suffering makes them ask frequently to stop the pain. They would prefer being dead. This contribution examines this kind of demand to find a helpful position for the caregiver and the patient. We should consider that near-death patients may often be at the climax of anxiety and depression is likely to switch over to dementia. In asking to finish with life, this purpose may change one day to another – sometimes it just highlights the wish to see things changed. The position of society and the medical staff has a high influence as well. Asking for euthanasia shows the variety of the same words that have different meanings from a medical, psychological, or psychopathological viewpoint. The purpose is to consider these aspects with the patient's demand.

Review Article **Published Date:-2024-01-03 15:06:01**

[Systemic Lupus Erythematosus and Depression Overview](#)

Systemic lupus erythematosus (SLE) is a chronic inflammatory multisystem disorder that commonly affects females during their reproductive years. It is characterized by the presence of autoantibodies and immune complex deposition, the etiology is not known but the interaction of an environmental agent in a genetically susceptible individual is thought to be fundamental. SLE most frequently involves the skin, joints, lungs, heart, kidney, and neuropsychiatric manifestations that may occur during the course of the disease. Mood disorders among SLE patients, particularly depression, are common and important psychiatric manifestations of the disease, in addition to their high incidence and possible deleterious influence on disease progression, so early identification and treatment of depression may have a significant influence on the patient's quality of life.
